

Drug Policy

Policy:	Spevigo® (spesolimab-sbzo subcutaneous injection - Boehringer Ingelheim)	Annual Review Date: 05/16/2024 Last Revised Date: 05/16/2024
----------------	---	---

OVERVIEW

Spevigo, an interleukin-36 receptor antagonist, is indicated for the treatment of generalized pustular psoriasis flares in adults and pediatric patients ≥ 12 years of age and weighing ≥ 40 kg.¹

Spevigo subcutaneous is used for treatment of generalized pustular psoriasis when patient is not experiencing a flare. The recommended dosage of Spevigo subcutaneous for treatment of generalized pustular psoriasis when not experiencing a flare in adults and pediatric patients ≥ 12 years of age and ≥ 40 kg is a loading dose of 600 mg (four 150 mg injections) followed by 300 mg (two 150 mg injections) administered subcutaneously 4 weeks later and every 4 weeks thereafter.

POLICY STATEMENT

This policy involves the use of Spevigo. Prior authorization is recommended for pharmacy benefit coverage of Spevigo. Approval is recommended for those who meet the conditions of coverage in the **Criteria and Initial/Extended Approval** for the diagnosis provided. **Conditions Not Recommended for Approval** are listed following the recommended authorization criteria. Requests for uses not listed in this policy will be reviewed for evidence of efficacy and for medical necessity on a case-by-case basis.

Because of the specialized skills required for evaluation and diagnosis of patients treated with Spevigo as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Spevigo be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals for initial therapy are provided for the initial approval duration noted below; if reauthorization is allowed, a response to therapy is required for continuation of therapy unless otherwise noted below.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Spevigo is recommended in those who meet the following criteria:

1. **Generalized Pustular Psoriasis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
 - A) **Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):
 - i. Patient is ≥ 12 years of age; AND
 - ii. Patient weighs ≥ 40 kilograms (kg); AND
 - iii. Patient has history of at least two generalized pustular psoriasis flares of moderate-to-severe intensity in the past; AND
 - iv. Patient has a Generalized Pustular Psoriasis Physician Global Assessment (GPPGA) total score of 0 or 1; AND
 - v. Patient meets ONE of the following (a or b):

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

Drug Policy

- a) Patient meets BOTH of the following ([1] and [2]):
 - (1) Patient has had a 4-month trial of least one treatment for generalized pustular psoriasis; AND
Note: Examples of treatment include methotrexate, acitretin, cyclosporine, or biologics.
 - (2) Patient has had a history of flaring while on treatment or with dose reduction or discontinuation of treatment; OR
 - b) Patient has tried at least one treatment for generalized pustular psoriasis but was unable to tolerate a 4-month trial; AND
 - vi. The medication is prescribed by or in consultation with a dermatologist.
- B) Patient is Currently Receiving Spevigo Subcutaneous.** Approve for 6 months if the patient meets BOTH of the following (i and ii):
- i. Patient has been established on therapy for at least 6 months; AND
Note: A patient who has received < 6 months of therapy or who is restarting therapy should be considered under criterion A (Initial Therapy).
 - ii. Patient has experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating the requested drug) in at least one of the following: reduction of generalized pustular psoriasis flares or an improvement in Generalized Pustular Psoriasis Physician Global Assessment (GPPGA) score.

Initial Approval/ Extended Approval.

A) Initial Approval: 6 months

B) Extended Approval: 6 months

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Spevigo has not been shown to be effective, or there are limited or preliminary data or potential safety concerns that are not supportive of general approval for the following conditions. (Note: This is not an exhaustive list of Conditions Not Recommended for Approval).

- 1. Concomitant use with Another Biologic or Disease-Modifying Antirheumatic Drugs (DMARD) Prescribed for Treatment of Generalized Pustular Psoriasis.** Although not approved, there are case reports documenting use of some biologics approved for plaque psoriasis (see [Appendix](#) for examples) for treatment of generalized pustular psoriasis. In the pivotal study, patients were required to discontinue therapy for generalized pustular psoriasis prior to receiving Spevigo.
Note: Patients with concomitant plaque psoriasis and generalized pustular psoriasis may be receiving a biologic for treatment of plaque psoriasis.
- 2. Plaque Psoriasis.** Spevigo has not been studied in patients with plaque psoriasis without generalized pustular psoriasis.
Note: Patients with concomitant plaque psoriasis and generalized pustular psoriasis may be reviewed under the generalized pustular psoriasis criteria above.
- 3. Synovitis-acne-pustulosis-hyperostosis-osteitis (SAPHO) syndrome.**

Drug Policy

4. **Primary erythrodermic psoriasis vulgaris.**
5. **Active infection, including clinically important localized infections.**
6. **Receipt of live vaccines (viral and/or bacterial) during therapy.**
7. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

Documentation Requirements:

The Company reserves the right to request additional documentation as part of its coverage determination process. The Company may deny reimbursement when it has determined that the drug provided or services performed were not medically necessary, investigational, or experimental, not within the scope of benefits afforded to the member and/or a pattern of billing or other practice has been found to be either inappropriate or excessive. Additional documentation supporting medical necessity for the services provided must be made available upon request to the Company. Documentation requested may include patient records, test results and/or credentials of the provider ordering or performing a service. The Company also reserves the right to modify, revise, change, apply and interpret this policy at its sole discretion, and the exercise of this discretion shall be final and binding.

REFERENCES

1. Spevigo [package insert]. Ridgefield, CT; Boehringer Ingelheim Pharmaceuticals, Inc.; March 2024. Accessed March 2024.
2. Bachelez H, Choon SE, Marrakchi S, et al; Effisayil 1 Trial Investigators. Trial of Spesolimab for Generalized Pustular Psoriasis. *N Engl J Med*. 2021 Dec 23;385(26):2431-2440. doi: 10.1056/NEJMoa2111563.
3. Choon SE, Lebwohl MG, Marrakchi S, et al. Study protocol of the global Effisayil 1 Phase II, multicentre, randomised, double-blind, placebo-controlled trial of spesolimab in patients with generalized pustular psoriasis presenting with an acute flare. *BMJ Open*. 2021 Mar 30;11(3):e043666. doi: 10.1136/bmjopen-2020-043666.
4. Navarini AA, Burden AD, Capon F, et al. European consensus statement on phenotypes of pustular psoriasis. *J Eur Acad Dermatol Venereol*. 2017 Nov;31(11):1792–1799. Crossref. PubMed. ISI.
5. Fujita H, Terui T, Hayama K, et al. Japanese guidelines for the management and treatment of generalized pustular psoriasis: the new pathogenesis and treatment of GPP. *J Dermatol*. 2018 Nov;45(11):1235–1270. Crossref. PubMed. ISI.
6. Morita A, Choon SE, Bachelez H, et al. Design of Effisayil™ 2: A Randomized, Double-Blind, Placebo-Controlled Study of Spesolimab in Preventing Flares in Patients with Generalized Pustular Psoriasis. *Dermatol Ther (Heidelb)*. 2023 Jan;13(1):347-359. doi: 10.1007/s13555-022-00835-6. Epub 2022 Nov 5. PMID: 36333618; PMCID: PMC9823166.
7. Burden AD, Bachelez H, Choon SE, et al. The Generalized Pustular Psoriasis Physician Global Assessment (GPPGA) score: online assessment and validation study of a specific measure of GPP disease activity, *British Journal of Dermatology*, Volume 189, Issue 1, July 2023, Pages 138–140, <https://doi.org/10.1093/bjd/ljad071>.
8. Robinson A, Van Voorhees AS, Hsu S, et al. Treatment of pustular psoriasis: from the medical board of the National Psoriasis Foundation. *J Am Acad Dermatol*. 2012;67(2):279-288.

APPENDIX

	Mechanism of Action	Examples of Inflammatory Indications*
Biologics		

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

Drug Policy

Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
Simponi®, Simponi® Aria™ (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
Actemra® (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PsA, RA
		IV formulation: JIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA [^] , RA
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC
Siliq™ (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection)	Inhibition of IL-17A	AS, ERA, nr-axSpA, PsO, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Ilumya™ (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
Skyrizi® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PsA, PsO
		IV formulation: CD
Tremfya™ (guselkumab SC injection)	Inhibition of IL-23	PsO
Entyvio™ (vedolizumab IV infusion)	Integrin receptor antagonist	CD, UC

* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; [^] Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis.